

ECE 302: Lecture 5.6 Sum of Two Random Variables

Prof Stanley Chan

School of Electrical and Computer Engineering
Purdue University



Sum of Two Random Variables

Given two random variables X and Y , what is the PDF of the sum, i.e., $X + Y$?

- $f_X(t) + f_Y(t)$?
- $f_X(t) \cdot f_Y(t)$?
- $f_X(x + y)$?
- ???

Outline

- Joint PDF and CDF
- Joint Expectation
- Conditional Distribution
- Conditional Expectation
- Sum of Two Random Variables
- Random Vectors
- High-dimensional Gaussians and Transformation
- Principal Component Analysis

Today's lecture

- Intuition
- Basic principle

Intuition

- $X =$ uniform in $0, 1, 2, 3$
- $Y =$ uniform in $0, 1, 2, 3$
- $Z = X + Y$

Find PMF of Z :

$$\begin{aligned}p_Z(0) &= \mathbb{P}[X + Y = 0] \\ &= \mathbb{P}[(X, Y) = (0, 0)] \\ &= p_X(0)p_Y(0) = \frac{1}{16}\end{aligned}$$

$$\begin{aligned}p_Z(1) &= \mathbb{P}[X + Y = 1] \\ &= \mathbb{P}[(X, Y) = (0, 1) \cup (1, 0)] \\ &= p_X(0)p_Y(1) + p_X(1)p_Y(0) = \frac{2}{16}.\end{aligned}$$

Intuition

$Z = X + Y$	Cases, written in terms of (X, Y)	Probability
0	(0,0)	1/16
1	(0,1), (1,0)	2/16
2	(1,1), (2,0), (0,2)	3/16
3	(3,0), (2,1), (1,2), (0,3)	4/16
4	(3,1), (2,2), (1,3)	3/16
5	(3,2), (2,3)	2/16
6	(3,3)	1/16

The shape of $X + Y$

Main result

Theorem

Let X and Y be two independent random variables with PDFs $f_X(x)$ and $f_Y(y)$ respectively. Let $Z = X + Y$. The PDF of Z is given by

$$f_Z(z) = (f_X * f_Y)(z) = \int_{-\infty}^{\infty} f_X(z - y)f_Y(y)dy, \quad (1)$$

where “*” denotes the convolution.

The PDF of $X + Y$ is the convolution $f_X * f_Y$.




Figure: When summing two random variables X and Y , we are effectively taking the convolution of the two respective PMF / PDFs.

Proof

Always start from CDF:

$$\begin{aligned}F_Z(z) &= \mathbb{P}[Z \leq z] \\ &= \mathbb{P}[X + Y \leq z].\end{aligned}$$

Now need to integrate:




Figure: The shaded region highlights the set $X + Y \leq Z$.

Proof

$$\begin{aligned}\mathbb{P}[X + Y \leq z] &= \int_{-\infty}^{\infty} \int_{-\infty}^{z-y} f_{X,Y}(x, y) dx dy \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{z-y} f_X(x) f_Y(y) dx dy, \quad (\text{independence})\end{aligned}$$


Finally, by fundamental theorem of calculus,

$$\begin{aligned}f_Z(z) &= \frac{d}{dz} F_Z(z) \\ &= \frac{d}{dz} \int_{-\infty}^{\infty} \int_{-\infty}^{z-y} f_X(x) f_Y(y) dx dy \\ &= \int_{-\infty}^{\infty} \left(\frac{d}{dz} \int_{-\infty}^{z-y} f_X(x) f_Y(y) dx \right) dy \\ &= \int_{-\infty}^{\infty} f_X(z-y) f_Y(y) dy = (f_X * f_Y)(z),\end{aligned}$$


Summary

How is convolution related to random variables?


- If you sum X and Y , the resulting PDF is the convolution of f_X and f_Y
- E.g., Convoluting two uniform random variables give you a triangle PDF.




(a) X_1



(b) $X_1 + X_2$



(c) $X_1 + \dots + X_5$



(d) $X_1 + \dots + X_{100}$

Questions?