

# Lecture 38

## Quantum Theory of Light

### 38.1 Quantum Theory of Light

#### 38.1.1 Historical Background

Quantum theory is a major intellectual achievement of the twentieth century, even though we are still discovering new knowledge in it. Several major experimental findings led to the revelation of quantum theory or quantum mechanics of nature. In nature, we know that many things are not infinitely divisible. Matter is not infinitely divisible as vindicated by the atomic theory of John Dalton (1766-1844) [221]. So fluid is not infinitely divisible: as when water is divided into smaller pieces, we will eventually arrive at water molecule,  $\text{H}_2\text{O}$ , which is the fundamental building block of water.

It turns out that electromagnetic energy is not infinitely divisible either. The electromagnetic radiation out of a heated cavity would obey a very different spectrum if electromagnetic energy is infinitely divisible. In order to fit experimental observation of radiation from a heated electromagnetic cavity, Max Planck (1900s) [222] proposed that electromagnetic energy comes in packets or is quantized. Each packet of energy or a quantum of energy  $E$  is associated with the frequency of electromagnetic wave, namely

$$E = \hbar\omega = \hbar 2\pi f = hf \quad (38.1.1)$$

where  $\hbar$  is now known as the Planck constant and  $\hbar = h/2\pi = 6.626 \times 10^{-34}$  J·s (Joule-second). Since  $\hbar$  is very small, this packet of energy is very small unless  $\omega$  is large. So it is no surprise that the quantization of electromagnetic field is first associated with light, a very high frequency electromagnetic radiation. A red-light photon at a wavelength of 700 nm corresponds to an energy of approximately  $2 \text{ eV} \approx 3 \times 10^{-19} \text{ J} \approx 75 k_B T$ , where  $k_B T$  denotes the thermal energy. A microwave photon has approximately  $1 \times 10^{-5} \text{ eV}$ .

The second experimental evidence that light is quantized is the photo-electric effect [223]. It was found that matter emitted electrons when light shined on it. First, the light frequency has to correspond to the “resonant” frequency of the atom. Second, the number of electrons emitted is proportional to the number of packets of energy  $\hbar\omega$  that the light carries. This

was a clear indication that light energy traveled in packets or quanta as posited by Einstein in 1905.

That light is a wave has been demonstrated by Newton's ring effect [224] in the eighteenth century (1717) (see Figure 38.1). In 1801, Thomas Young demonstrated the double slit experiment [225] that further confirmed the wave nature of light (see Figure 38.2). But by the beginning of the 20-th century, one has to accept that light is both a particle, called a photon, carrying a quantum of energy with momentum, as well as a particle endowed with wave-like behavior. This is called wave-particle duality.

**Theory**




Fig. 1 Experimental set-up to observe Newton's ring




Fig. 2. Newton's rings

Figure 38.1: A Newton's rings experiment (courtesy of [224]).




Figure 38.2: A Young's double-slit experiment (courtesy of [226]).

This concept was not new to quantum theory as electrons were known to behave both like a particle and a wave. The particle nature of an electron was confirmed by the measurement of its charge by Millikan in 1913 in his oil-drop experiment. (The double slit experiment for electron was done in 1927 by Davison and Germer, indicating that an electron has a wave nature [225].) In 1924, De Broglie [227] suggested that there is a wave associated with an electron with momentum  $p$  such that

$$p = \hbar k \quad (38.1.2)$$

where  $k = 2\pi/\lambda$ , the wavenumber. All this knowledge gave hint to the quantum theorists of that era to come up with a new way to describe nature.

Classically, particles like an electron moves through space obeying Newton's laws of motion first established in 1687 [228]. Old ways of describing particle motion were known as classical mechanics, and the new way of describing particle motion is known as quantum mechanics. Quantum mechanics is very much motivated by a branch of classical mechanics called Hamiltonian mechanics. We will first use Hamiltonian mechanics to study a simple pendulum and connect it with electromagnetic oscillations.

### 38.1.2 Connecting Electromagnetic Oscillation to Simple Pendulum

The quantization of electromagnetic field theory was started by Dirac in 1927 [3]. In the beginning, it was called quantum electrodynamics important for understanding particle physics

phenomena and light-matter interactions [229]. Later on, it became important in quantum optics where quantum effects in electromagnetics technologies first emerged. Now, microwave photons are measurable and are important in quantum computers. Hence, quantum effects are important in the microwave regime as well.

Maxwell's equations can be regarded as for describing an infinite set of coupled harmonic oscillators. In one dimension, when a wave propagates on a string, or an electromagnetic wave propagating on a transmission line, they can be regarded as propagating on a set of coupled harmonic oscillators as shown in Figure 38.3. Maxwell's equations describe the waves travelling in 3D space due to the coupling between an infinite set of harmonic oscillators. In fact, methods have been developed to solve Maxwell's equations using transmission-line-matrix (TLM) method [230], or the partial element equivalent circuit (PEEC) method [231].




Figure 38.3: Maxwell's equations describe the coupling of harmonic oscillators in a 3D space. This is similar to waves propagating on a string or a 1D transmission line, or a 2D array of coupled oscillators. The saw-tooth symbol in the figure represents a spring.

The cavity modes in electromagnetics are similar to the oscillation of a pendulum. To understand the quantization of electromagnetic field, we start by connecting these cavity modes oscillations to the oscillations of a simple pendulum. It is to be noted that fundamentally, electromagnetic oscillation exists because of displacement current. Displacement current exists even in vacuum because vacuum is polarizable, namely that  $\mathbf{D} = \epsilon\mathbf{E}$ . Furthermore, displacement current exists because of the  $\partial\mathbf{D}/\partial t$  term in the generalized Ampere's law added by Maxwell, namely,

$$\nabla \times \mathbf{H} = \frac{\partial \mathbf{D}}{\partial t} + \mathbf{J} \quad (38.1.3)$$

Together with Faraday’s law that

$$\nabla \times \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \tag{38.1.4}$$

(38.1.3) and (38.1.4) together allow for the existence of wave. The coupling between the two equations gives rise to the “springiness” of electromagnetic fields.

Wave exists due to the existence of coupled harmonic oscillators, and at a fundamental level, these harmonic oscillators are electron-positron (e-p) pairs. The fact that they are coupled allows waves to propagate through space, and even in vacuum.




Figure 38.4: A one-dimensional cavity solution to Maxwell’s equations is one of the simplest way to solve Maxwell’s equations.

To make life simpler, we can start by looking at a one dimensional cavity formed by two PEC (perfect electric conductor) plates as shown in Figure 38.4. Assume source-free Maxwell’s equations in between the plates and letting  $\mathbf{E} = \hat{x}E_x$ ,  $\mathbf{H} = \hat{y}H_y$ , then (38.1.3) and (38.1.4) become

$$\frac{\partial}{\partial z} H_y = -\epsilon \frac{\partial}{\partial t} E_x \tag{38.1.5}$$

$$\frac{\partial}{\partial z} E_x = -\mu \frac{\partial}{\partial t} H_y \tag{38.1.6}$$

The above are similar to the telegrapher’s equations. We can combine them to arrive at

$$\frac{\partial^2}{\partial z^2} E_x = \mu\epsilon \frac{\partial^2}{\partial t^2} E_x \tag{38.1.7}$$

There are infinitely many ways to solve the above partial differential equation. But here, we use separation of variables to solve the above by letting  $E_x(z, t) = E_0(t)f(z)$ . Then we arrive

at two separate equations that

$$\frac{d^2 E_0(t)}{dt^2} = -\omega_l^2 E_0(t) \quad (38.1.8)$$

and

$$\frac{d^2 f(z)}{dz^2} = -\omega_l^2 \mu \epsilon f(z) \quad (38.1.9)$$

where  $\omega_l^2$  is the separation constant. There are infinitely many ways to solve the above equations which are also eigenvalue equations where  $\omega_l^2$  and  $\omega_l^2 \mu \epsilon$  are eigenvalues for the first and the second equations, respectively. The general solution for (38.1.9) is that

$$E_0(t) = E_0 \cos(\omega_l t + \psi) \quad (38.1.10)$$

In the above,  $\omega_l$ , which is related to the separation constant, is yet indeterminate. To make  $\omega_l^2$  determinate, we need to impose boundary conditions. A simple way is to impose homogeneous Dirichlet boundary conditions that  $f(z) = 0$  at  $z = 0$  and  $z = L$ . This implies that  $f(z) = \sin(k_l z)$ . In order to satisfy the boundary conditions at  $z = 0$  and  $z = L$ , one deduces that

$$k_l = \frac{l\pi}{L}, \quad l = 1, 2, 3, \dots \quad (38.1.11)$$

Then,

$$\frac{\partial^2 f(z)}{\partial z^2} = -k_l^2 f(z) \quad (38.1.12)$$

where  $k_l^2 = \omega_l^2 \mu \epsilon$ . Hence,  $k_l = \omega_l/c$ , and the above solution can only exist for discrete frequencies or that

$$\omega_l = \frac{l\pi}{L} c, \quad l = 1, 2, 3, \dots \quad (38.1.13)$$

These are the discrete resonant frequencies  $\omega_l$  of the modes of the 1D cavity.

The above solutions for  $E_x(z, t)$  can be thought of as the collective oscillations of coupled harmonic oscillators forming the modes of the cavity. At the fundamental level, these oscillations are oscillators made by electron-positron pairs. But macroscopically, their collective resonances manifest themselves as giving rise to infinitely many electromagnetic cavity modes. The amplitudes of these modes,  $E_0(t)$  are simple harmonic oscillations.

The resonance between two parallel PEC plates is similar to the resonance of a transmission line of length  $L$  shorted at both ends. One can see that the resonance of a shorted transmission line is similar to the coupling of infinitely many LC tank circuits. To see this, as shown in Figure 38.3, we start with a single LC tank circuit as a simple harmonic oscillator with only one resonant frequency. When two LC tank circuits are coupled to each other, they will have two resonant frequencies. For  $N$  of them, they will have  $N$  resonant frequencies. For a continuum of them, they will be infinitely many resonant frequencies or modes as indicated by (38.1.11).

What is more important is that the resonance of each of these modes is similar or homomorphic to the resonance of a simple pendulum or a simple harmonic oscillator. For a fixed point in space, the field due to this oscillation is similar to the oscillation of a simple pendulum.

As we have seen in the Drude-Lorentz-Sommerfeld mode, for a particle of mass  $m$  attached to a spring connected to a wall, where the restoring force is like Hooke's law, the equation of motion of a pendulum by Newton's law is

$$m \frac{d^2 x}{dt^2} + \kappa x = 0 \quad (38.1.14)$$

where  $\kappa$  is the spring constant, and we assume that the oscillator is not driven by an external force, but is in natural or free oscillation. By letting<sup>1</sup>

$$x = x_0 e^{-i\omega t} \quad (38.1.15)$$

the above becomes

$$-m\omega^2 x_0 + \kappa x_0 = 0 \quad (38.1.16)$$

Again, a non-trivial solution is possible only at the resonant frequency of the oscillator or that when  $\omega = \omega_0$  where

$$\omega_0 = \sqrt{\frac{\kappa}{m}} \quad (38.1.17)$$

## 38.2 Hamiltonian Mechanics

Equation (38.1.14) can be derived by Newton's law but it can also be derived via Hamiltonian mechanics. Since Hamiltonian mechanics motivates quantum mechanics, we will look at the Hamiltonian mechanics view of the equation of motion (EOM) of a simple pendulum given by (38.1.14).

Hamiltonian mechanics, developed by Hamilton (1805-1865) [232], is motivated by energy conservation. The Hamiltonian  $H$  of a system is given by its total energy, namely that

$$H = T + V \quad (38.2.1)$$

where  $T$  is the kinetic energy and  $V$  is the potential energy of the system.

For a simple pendulum, the kinetic energy is given by

$$T = \frac{1}{2} m v^2 = \frac{1}{2m} m^2 v^2 = \frac{p^2}{2m} \quad (38.2.2)$$

where  $p = mv$  is the momentum of the particle. The potential energy, assuming that the particle is attached to a spring with spring constant  $\kappa$ , is given by

$$V = \frac{1}{2} \kappa x^2 = \frac{1}{2} m \omega_0^2 x^2 \quad (38.2.3)$$

---

<sup>1</sup>For this part of the lecture, we will switch to using  $\exp(-i\omega t)$  time convention as is commonly used in optics and physics literatures.

Hence, the Hamiltonian is given by

$$H = T + V = \frac{p^2}{2m} + \frac{1}{2}m\omega_0^2x^2 \quad (38.2.4)$$

At any instant of time  $t$ , we assume that  $p(t) = mv(t) = m\frac{d}{dt}x(t)$  is independent of  $x(t)$ .<sup>2</sup> In other words, they can vary independently of each other. But  $p(t)$  and  $x(t)$  have to time evolve to conserve energy to keep  $H$ , the total energy, constant or independent of time. In other words,

$$\frac{d}{dt}H[p(t), x(t)] = 0 = \frac{dp}{dt}\frac{\partial H}{\partial p} + \frac{dx}{dt}\frac{\partial H}{\partial x} \quad (38.2.5)$$

Therefore, the Hamilton equations of motion are derived to be<sup>3</sup>

$$\frac{dp}{dt} = -\frac{\partial H}{\partial x}, \quad \frac{dx}{dt} = \frac{\partial H}{\partial p} \quad (38.2.6)$$

From (38.2.4), we gather that

$$\frac{\partial H}{\partial x} = m\omega_0^2x, \quad \frac{\partial H}{\partial p} = \frac{p}{m} \quad (38.2.7)$$

Applying (38.2.6), we have<sup>4</sup>

$$\frac{dx}{dt} = \frac{p}{m}, \quad \frac{dp}{dt} = -m\omega_0^2x \quad (38.2.8)$$

Combining the two equations in (38.2.8) above, we have

$$m\frac{d^2x}{dt^2} = -m\omega_0^2x = -\kappa x \quad (38.2.9)$$

which is also derivable by Newton's law.

A typical harmonic oscillator solution to (38.2.9) is

$$x(t) = x_0 \cos(\omega_0 t + \psi) \quad (38.2.10)$$

The corresponding  $p(t) = m\frac{dx}{dt}$  is

$$p(t) = -mx_0\omega_0 \sin(\omega_0 t + \psi) \quad (38.2.11)$$

Hence

$$\begin{aligned} H &= \frac{1}{2}m\omega_0^2x_0^2 \sin^2(\omega_0 t + \psi) + \frac{1}{2}m\omega_0^2x_0^2 \cos^2(\omega_0 t + \psi) \\ &= \frac{1}{2}m\omega_0^2x_0^2 = E \end{aligned} \quad (38.2.12)$$

And the total energy  $E$  very much depends on the amplitude  $x_0$  of the oscillation.

<sup>2</sup> $p(t)$  and  $x(t)$  are termed conjugate variables in many textbooks.

<sup>3</sup>Note that the Hamilton equations are determined to within a multiplicative constant, because one has not stipulated the connection between space and time, or we have not calibrated our clock [233].

<sup>4</sup>We can also calibrate our clock here so that it agrees with our definition of momentum in the ensuing equation.



### 38.3 Schrodinger Equation (1925)

Having seen the Hamiltonian mechanics for describing a simple pendulum which is homomorphic to a cavity resonator, we shall next see the quantum mechanics description of the same simple pendulum: In other words, we will look at a quantum pendulum. To this end, we will invoke Schrodinger equation.

Schrodinger equation cannot be derived just as in the case Maxwell's equations. It is a wonderful result of a postulate and a guessing game based on experimental observations [63,64]. Hamiltonian mechanics says that

$$H = \frac{p^2}{2m} + \frac{1}{2}m\omega_0^2x^2 = E \quad (38.3.1)$$

where  $E$  is the total energy of the oscillator, or pendulum. In classical mechanics, the position  $x$  of the particle associated with the pendulum is known with great certainty. But in the quantum world, this position  $x$  of the quantum particle is uncertain and is fuzzy.

To build this uncertainty into a quantum harmonic oscillator, we have to look at it from the quantum world. The position of the particle is described by a wave function,<sup>5</sup> which makes the location of the particle uncertain. To this end, Schrodinger proposed his equation which is a partial differential equation. He was very much motivated by the experimental revelation then that  $p = \hbar k$  from De Broglie and that  $E = \hbar\omega$  from Planck's law. Equation (38.3.1) can be written more suggestively as

$$\frac{\hbar^2k^2}{2m} + \frac{1}{2}m\omega_0^2x^2 = \hbar\omega \quad (38.3.2)$$

To add more depth to the above equation, one lets the above become an operator equation that operates on a wave function  $\psi(x, t)$  so that

$$-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} \psi(x, t) + \frac{1}{2}m\omega_0^2x^2\psi(x, t) = i\hbar \frac{\partial}{\partial t} \psi(x, t) \quad (38.3.3)$$

If the wave function is of the form

$$\psi(x, t) \sim e^{ikx - i\omega t} \quad (38.3.4)$$

then upon substituting (38.3.4) into (38.3.3), we retrieve (38.3.2).

Equation (38.3.3) is Schrodinger equation in one dimension for the quantum version of the simple harmonic oscillator. In Schrodinger equation, we can further posit that the wave function has the general form

$$\psi(x, t) = e^{ikx - i\omega t} A(x, t) \quad (38.3.5)$$

where  $A(x, t)$  is a slowly varying function of  $x$  and  $t$ , compared to  $e^{ikx - i\omega t}$ .<sup>6</sup> In other words, this is the expression for a wave packet. With this wave packet, the  $\partial^2/\partial x^2$  can be again

<sup>5</sup>Since a function is equivalent to a vector, and this wave function describes the state of the quantum system, this is also called a state vector.

<sup>6</sup>This is similar in spirit when we study high frequency solutions of Maxwell's equations and paraxial wave approximation.

approximated by  $-k^2$  as has been done in the paraxial wave approximation. Furthermore, if the signal is assumed to be quasi-monochromatic, then  $i\hbar\partial/\partial t\psi(x, t) \approx \hbar\omega$ , we again retrieve the classical equation in (38.3.2) from (38.3.3). Hence, the classical equation (38.3.2) is a short wavelength, monochromatic approximation of Schrodinger equation. However, as we shall see, the solutions to Schrodinger equation are not limited to just wave packets described by (38.3.5).

For this course, we need only to study the one-dimensional Schrodinger equation. The above can be converted into eigenvalue problem, just as in waveguide and cavity problems, using separation of variables, by letting<sup>7</sup>

$$\psi(x, t) = \psi_n(x)e^{-i\omega_n t} \quad (38.3.6)$$

By so doing, (38.3.3) becomes

$$\left[ -\frac{\hbar^2}{2m} \frac{d^2}{dx^2} + \frac{1}{2}m\omega_0^2 x^2 \right] \psi_n(x) = E_n \psi_n(x) \quad (38.3.7)$$

where  $E_n = \hbar\omega_n$  is the eigenvalue of the problem while  $\psi_n(x)$  is the eigenfunction.

The parabolic  $x^2$  potential profile is also known as a potential well as it can provide the restoring force to keep the particle bound to the well classically. The above equation is also similar to the electromagnetic equation for a dielectric slab waveguide, where the second term is a dielectric profile (mind you, varying in the  $x$  direction) that can trap a waveguide mode. Therefore, the potential well is a trap for the particle both in classical mechanics or wave physics.

The above equation (38.3.7) can be solved in closed form in terms of Hermite-Gaussian functions (1864) [234], or that

$$\psi_n(x) = \sqrt{\frac{1}{2^n n!}} \sqrt{\frac{m\omega_0}{\pi\hbar}} e^{-\frac{m\omega_0}{2\hbar} x^2} H_n \left( \sqrt{\frac{m\omega_0}{\hbar}} x \right) \quad (38.3.8)$$

where  $H_n(y)$  is a Hermite polynomial, and the eigenvalues are

$$E_n = \left( n + \frac{1}{2} \right) \hbar\omega_0 \quad (38.3.9)$$

Here, the eigenfunction or eigenstate  $\psi_n(x)$  is known as the photon number state (or just a number state) of the solution. It corresponds to having  $n$  photons in the oscillation. If this is conceived as the collective oscillation of the e-p pairs in a cavity, there are  $n$  photons corresponding to energy of  $n\hbar\omega_0$  embedded in the collective oscillation. The larger  $E_n$  is, the larger the number of photons there is. However, there is a curious mode at  $n = 0$ . This corresponds to no photon, and yet, there is a wave function  $\psi_0(x)$ . This is the zero-point energy state. This state is there even if the system is at its lowest energy state.

It is to be noted that in the quantum world, the position  $x$  of the pendulum is random. Moreover, this position  $x(t)$  is mapped to the amplitude  $E_0(t)$  of the field. Hence, it is the

<sup>7</sup>Mind you, the following is  $\omega_n$ , not  $\omega_0$ .

amplitude of an electromagnetic oscillation that becomes uncertain and fuzzy as shown in Figure 38.5.




Figure 38.5: Schematic representation of the randomness of measured electric field. The electric field amplitude maps to the displacement (position) of the quantum harmonic oscillator, which is a random variable (courtesy of Kira and Koch [235]).




Figure 38.6: Plots of the eigensolutions of the quantum harmonic oscillator (courtesy of Wiki [236]).

### 38.4 Some Quantum Interpretations—A Preview

Schrodinger used this equation with resounding success. He derived a three-dimensional version of this to study the wave function and eigenvalues of a hydrogen atom. These eigenvalues  $E_n$  for a hydrogen atom agreed well with experimental observations that had eluded scientists for decades. Schrodinger did not actually understand what this wave function meant. It was Max Born (1926) who gave a physical interpretation of this wave function.

Given a wave function  $\psi(x, t)$ , then  $|\psi(x, t)|^2 \Delta x$  is the probability of finding the particle in the interval  $[x, x + \Delta x]$ . Therefore,  $|\psi(x, t)|^2$  is a probability density function (PDF), and it is necessary that

$$\int_{-\infty}^{\infty} dx |\psi(x, t)|^2 = 1 \tag{38.4.1}$$

The position  $x$  of the particle is uncertain and is now a random variable. The average value or expectation value of  $x$  is given by

$$\int_{-\infty}^{\infty} dx x |\psi(x, t)|^2 = \langle x(t) \rangle = \bar{x}(t) \tag{38.4.2}$$

This is not the most ideal notation, since although  $x$  is not a function of time, its expectation value with respect to a time-varying function,  $\psi(x, t)$ , can be time-varying.

Notice that in going from (38.3.1) to (38.3.3), or from a classical picture to a quantum picture, we have let the momentum become  $p$ , originally a scalar number in the classical world, become a differential operator, namely that

$$p \rightarrow \hat{p} = -i\hbar \frac{\partial}{\partial x} \quad (38.4.3)$$

The momentum of a particle also becomes uncertain, and its expectation value is given by

$$\int_{-\infty}^{\infty} dx \psi^*(x, t) \hat{p} \psi(x, t) = -i\hbar \int_{-\infty}^{\infty} dx \psi^*(x, t) \frac{\partial}{\partial x} \psi(x, t) = \langle \hat{p}(t) \rangle = \bar{p}(t) \quad (38.4.4)$$

The expectation values of position  $x$  and the momentum operator  $\hat{p}$  are measurable in the laboratory. Hence, they are also called observables.

One more very important aspect of quantum theory is that since  $p \rightarrow \hat{p} = -i\hbar \frac{\partial}{\partial x}$ ,  $\hat{p}$  and  $x$  do not commute. In other words, it can be shown that

$$[\hat{p}, x] = \left[ -i\hbar \frac{\partial}{\partial x}, x \right] = -i\hbar \quad (38.4.5)$$

In the classical world,  $[p, x] = 0$ , but not in the quantum world. In the equation above, we can elevate  $x$  to become an operator by letting  $\hat{x} = x\hat{I}$ , where  $\hat{I}$  is the identity operator. Then both  $\hat{p}$  and  $\hat{x}$  are now operators, and are on the same footing. In this manner, we can rewrite equation (38.4.5) above as

$$[\hat{p}, \hat{x}] = -i\hbar \hat{I} \quad (38.4.6)$$

It can be shown easily that when two operators share the same set of eigenfunctions, they commute. When two operators  $\hat{p}$  and  $\hat{x}$  do not commute, it means that the expectation values of quantities associated with the operators,  $\langle \hat{p} \rangle$  and  $\langle \hat{x} \rangle$ , cannot be determined to arbitrary precision simultaneously. For instance,  $\hat{p}$  and  $\hat{x}$  correspond to random variables, then the standard deviation of their measurable values, or their expectation values, obey the uncertainty principle relationship that<sup>8</sup>

$$\Delta p \Delta x \geq \hbar/2 \quad (38.4.7)$$

where  $\Delta p$  and  $\Delta x$  are the standard deviation of the random variables  $p$  and  $x$ .

## 38.5 Bizarre Nature of the Photon Number States

The photon number states are successful in predicting that the collective e-p oscillations are associated with  $n$  photons embedded in the energy of the oscillating modes. However, these number states are bizarre: The expectation values of the position of the quantum pendulum associated these states are always zero. To illustrate further, we form the wave function with a photon-number state

$$\psi(x, t) = \psi_n(x) e^{-i\omega_n t}$$

<sup>8</sup>The proof of this is quite straightforward but is outside the scope of this course.

Previously, since the  $\psi_n(x)$  are eigenfunctions, they are mutually orthogonal and they can be orthonormalized meaning that

$$\int_{-\infty}^{\infty} dx \psi_n^*(x) \psi_{n'}(x) = \delta_{nn'} \quad (38.5.1)$$

Then one can easily show that the expectation value of the position of the quantum pendulum in a photon number state is

$$\langle x(t) \rangle = \bar{x}(t) = \int_{-\infty}^{\infty} dx x |\psi(x, t)|^2 = \int_{-\infty}^{\infty} dx x |\psi_n(x)|^2 = 0 \quad (38.5.2)$$

because the integrand is always odd symmetric. In other words, the expectation value of the position  $x$  of the pendulum is always zero. It can also be shown that the expectation value of the momentum operator  $\hat{p}$  is also zero for these photon number states. Hence, there are no classical oscillations that resemble them. Therefore, one has to form new wave functions by linear superposing these photon number states into a coherent state. This will be the discussion next.

# Bibliography

- [1] J. A. Kong, *Theory of electromagnetic waves*. New York, Wiley-Interscience, 1975.
- [2] A. Einstein *et al.*, “On the electrodynamics of moving bodies,” *Annalen der Physik*, vol. 17, no. 891, p. 50, 1905.
- [3] P. A. M. Dirac, “The quantum theory of the emission and absorption of radiation,” *Proceedings of the Royal Society of London. Series A, Containing Papers of a Mathematical and Physical Character*, vol. 114, no. 767, pp. 243–265, 1927.
- [4] R. J. Glauber, “Coherent and incoherent states of the radiation field,” *Physical Review*, vol. 131, no. 6, p. 2766, 1963.
- [5] C.-N. Yang and R. L. Mills, “Conservation of isotopic spin and isotopic gauge invariance,” *Physical review*, vol. 96, no. 1, p. 191, 1954.
- [6] G. t’Hooft, *50 years of Yang-Mills theory*. World Scientific, 2005.
- [7] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*. Princeton University Press, 2017.
- [8] F. Teixeira and W. C. Chew, “Differential forms, metrics, and the reflectionless absorption of electromagnetic waves,” *Journal of Electromagnetic Waves and Applications*, vol. 13, no. 5, pp. 665–686, 1999.
- [9] W. C. Chew, E. Michielssen, J.-M. Jin, and J. Song, *Fast and efficient algorithms in computational electromagnetics*. Artech House, Inc., 2001.
- [10] A. Volta, “On the electricity excited by the mere contact of conducting substances of different kinds. in a letter from Mr. Alexander Volta, FRS Professor of Natural Philosophy in the University of Pavia, to the Rt. Hon. Sir Joseph Banks, Bart. KBPR S,” *Philosophical transactions of the Royal Society of London*, no. 90, pp. 403–431, 1800.
- [11] A.-M. Ampère, *Exposé méthodique des phénomènes électro-dynamiques, et des lois de ces phénomènes*. Bachelier, 1823.
- [12] —, *Mémoire sur la théorie mathématique des phénomènes électro-dynamiques uniquement déduite de l’expérience: dans lequel se trouvent réunis les Mémoires que M. Ampère a communiqués à l’Académie royale des Sciences, dans les séances des 4 et*

26 décembre 1820, 10 juin 1822, 22 décembre 1823, 12 septembre et 21 novembre 1825. Bachelier, 1825.

- [13] B. Jones and M. Faraday, *The life and letters of Faraday*. Cambridge University Press, 2010, vol. 2.
- [14] G. Kirchhoff, “Ueber die auflösung der gleichungen, auf welche man bei der untersuchung der linearen vertheilung galvanischer ströme geführt wird,” *Annalen der Physik*, vol. 148, no. 12, pp. 497–508, 1847.
- [15] L. Weinberg, “Kirchhoff’s’ third and fourth laws’,” *IRE Transactions on Circuit Theory*, vol. 5, no. 1, pp. 8–30, 1958.
- [16] T. Standage, *The Victorian Internet: The remarkable story of the telegraph and the nineteenth century’s online pioneers*. Phoenix, 1998.
- [17] J. C. Maxwell, “A dynamical theory of the electromagnetic field,” *Philosophical transactions of the Royal Society of London*, no. 155, pp. 459–512, 1865.
- [18] H. Hertz, “On the finite velocity of propagation of electromagnetic actions,” *Electric Waves*, vol. 110, 1888.
- [19] M. Romer and I. B. Cohen, “Roemer and the first determination of the velocity of light (1676),” *Isis*, vol. 31, no. 2, pp. 327–379, 1940.
- [20] A. Arons and M. Peppard, “Einstein’s proposal of the photon concept—a translation of the Annalen der Physik paper of 1905,” *American Journal of Physics*, vol. 33, no. 5, pp. 367–374, 1965.
- [21] A. Pais, “Einstein and the quantum theory,” *Reviews of Modern Physics*, vol. 51, no. 4, p. 863, 1979.
- [22] M. Planck, “On the law of distribution of energy in the normal spectrum,” *Annalen der physik*, vol. 4, no. 553, p. 1, 1901.
- [23] Z. Peng, S. De Graaf, J. Tsai, and O. Astafiev, “Tuneable on-demand single-photon source in the microwave range,” *Nature communications*, vol. 7, p. 12588, 2016.
- [24] B. D. Gates, Q. Xu, M. Stewart, D. Ryan, C. G. Willson, and G. M. Whitesides, “New approaches to nanofabrication: molding, printing, and other techniques,” *Chemical reviews*, vol. 105, no. 4, pp. 1171–1196, 2005.
- [25] J. S. Bell, “The debate on the significance of his contributions to the foundations of quantum mechanics, Bells Theorem and the Foundations of Modern Physics (A. van der Merwe, F. Selleri, and G. Tarozzi, eds.),” 1992.
- [26] D. J. Griffiths and D. F. Schroeter, *Introduction to quantum mechanics*. Cambridge University Press, 2018.
- [27] C. Pickover, *Archimedes to Hawking: Laws of science and the great minds behind them*. Oxford University Press, 2008.



- [28] R. Resnick, J. Walker, and D. Halliday, *Fundamentals of physics*. John Wiley, 1988.
- [29] S. Ramo, J. R. Whinnery, and T. Duzer van, *Fields and waves in communication electronics, Third Edition*. John Wiley & Sons, Inc., 1995, also 1965, 1984.
- [30] J. L. De Lagrange, “Recherches d’arithmétique,” *Nouveaux Mémoires de l’Académie de Berlin*, 1773.
- [31] J. A. Kong, *Electromagnetic Wave Theory*. EMW Publishing, 2008, also 1985.
- [32] H. M. Schey, *Div, grad, curl, and all that: an informal text on vector calculus*. WW Norton New York, 2005.
- [33] R. P. Feynman, R. B. Leighton, and M. Sands, *The Feynman lectures on physics, Vols. I, II, & III: The new millennium edition*. Basic books, 2011, also 1963, 2006, vol. 1,2,3.
- [34] W. C. Chew, *Waves and fields in inhomogeneous media*. IEEE Press, 1995, also 1990.
- [35] V. J. Katz, “The history of Stokes’ theorem,” *Mathematics Magazine*, vol. 52, no. 3, pp. 146–156, 1979.
- [36] W. K. Panofsky and M. Phillips, *Classical electricity and magnetism*. Courier Corporation, 2005.
- [37] T. Lancaster and S. J. Blundell, *Quantum field theory for the gifted amateur*. OUP Oxford, 2014.
- [38] W. C. Chew, “Fields and waves: Lecture notes for ECE 350 at UIUC,” <https://engineering.purdue.edu/wcchew/ece350.html>, 1990.
- [39] C. M. Bender and S. A. Orszag, *Advanced mathematical methods for scientists and engineers I: Asymptotic methods and perturbation theory*. Springer Science & Business Media, 2013.
- [40] J. M. Crowley, *Fundamentals of applied electrostatics*. Krieger Publishing Company, 1986.
- [41] C. Balanis, *Advanced Engineering Electromagnetics*. Hoboken, NJ, USA: Wiley, 2012.
- [42] J. D. Jackson, *Classical electrodynamics*. John Wiley & Sons, 1999.
- [43] R. Courant and D. Hilbert, *Methods of Mathematical Physics, Volumes 1 and 2*. Interscience Publ., 1962.
- [44] L. Esaki and R. Tsu, “Superlattice and negative differential conductivity in semiconductors,” *IBM Journal of Research and Development*, vol. 14, no. 1, pp. 61–65, 1970.
- [45] E. Kudeki and D. C. Munson, *Analog Signals and Systems*. Upper Saddle River, NJ, USA: Pearson Prentice Hall, 2009.
- [46] A. V. Oppenheim and R. W. Schaffer, *Discrete-time signal processing*. Pearson Education, 2014.

- [47] R. F. Harrington, *Time-harmonic electromagnetic fields*. McGraw-Hill, 1961.
- [48] E. C. Jordan and K. G. Balmain, *Electromagnetic waves and radiating systems*. Prentice-Hall, 1968.
- [49] G. Agarwal, D. Pattanayak, and E. Wolf, “Electromagnetic fields in spatially dispersive media,” *Physical Review B*, vol. 10, no. 4, p. 1447, 1974.
- [50] S. L. Chuang, *Physics of photonic devices*. John Wiley & Sons, 2012, vol. 80.
- [51] B. E. Saleh and M. C. Teich, *Fundamentals of photonics*. John Wiley & Sons, 2019.
- [52] M. Born and E. Wolf, *Principles of optics: electromagnetic theory of propagation, interference and diffraction of light*. Elsevier, 2013, also 1959 to 1986.
- [53] R. W. Boyd, *Nonlinear optics*. Elsevier, 2003.
- [54] Y.-R. Shen, *The principles of nonlinear optics*. New York, Wiley-Interscience, 1984.
- [55] N. Bloembergen, *Nonlinear optics*. World Scientific, 1996.
- [56] P. C. Krause, O. Wasynczuk, and S. D. Sudhoff, *Analysis of electric machinery*. McGraw-Hill New York, 1986.
- [57] A. E. Fitzgerald, C. Kingsley, S. D. Umans, and B. James, *Electric machinery*. McGraw-Hill New York, 2003, vol. 5.
- [58] M. A. Brown and R. C. Semelka, *MRI.: Basic Principles and Applications*. John Wiley & Sons, 2011.
- [59] C. A. Balanis, *Advanced engineering electromagnetics*. John Wiley & Sons, 1999, also 1989.
- [60] Wikipedia, “Lorentz force,” [https://en.wikipedia.org/wiki/Lorentz\\_force/](https://en.wikipedia.org/wiki/Lorentz_force/), accessed: 2019-09-06.
- [61] R. O. Dendy, *Plasma physics: an introductory course*. Cambridge University Press, 1995.
- [62] P. Sen and W. C. Chew, “The frequency dependent dielectric and conductivity response of sedimentary rocks,” *Journal of microwave power*, vol. 18, no. 1, pp. 95–105, 1983.
- [63] D. A. Miller, *Quantum Mechanics for Scientists and Engineers*. Cambridge, UK: Cambridge University Press, 2008.
- [64] W. C. Chew, “Quantum mechanics made simple: Lecture notes for ECE 487 at UIUC,” <http://wcchew.ece.illinois.edu/chew/course/QMAll20161206.pdf>, 2016.
- [65] B. G. Streetman and S. Banerjee, *Solid state electronic devices*. Prentice hall Englewood Cliffs, NJ, 1995.

- [66] Smithsonian, “This 1600-year-old goblet shows that the romans were nanotechnology pioneers,” <https://www.smithsonianmag.com/history/this-1600-year-old-goblet-shows-that-the-romans-were-nanotechnology-pioneers-787224/>, accessed: 2019-09-06.
- [67] K. G. Budden, *Radio waves in the ionosphere*. Cambridge University Press, 2009.
- [68] R. Fitzpatrick, *Plasma physics: an introduction*. CRC Press, 2014.
- [69] G. Strang, *Introduction to linear algebra*. Wellesley-Cambridge Press Wellesley, MA, 1993, vol. 3.
- [70] K. C. Yeh and C.-H. Liu, “Radio wave scintillations in the ionosphere,” *Proceedings of the IEEE*, vol. 70, no. 4, pp. 324–360, 1982.
- [71] J. Kraus, *Electromagnetics*. McGraw-Hill, 1984, also 1953, 1973, 1981.
- [72] Wikipedia, “Circular polarization,” [https://en.wikipedia.org/wiki/Circular\\_polarization](https://en.wikipedia.org/wiki/Circular_polarization).
- [73] Q. Zhan, “Cylindrical vector beams: from mathematical concepts to applications,” *Advances in Optics and Photonics*, vol. 1, no. 1, pp. 1–57, 2009.
- [74] H. Haus, *Electromagnetic Noise and Quantum Optical Measurements*, ser. Advanced Texts in Physics. Springer Berlin Heidelberg, 2000.
- [75] W. C. Chew, “Lectures on theory of microwave and optical waveguides, for ECE 531 at UIUC,” <https://engineering.purdue.edu/wcchew/course/tqwAll20160215.pdf>, 2016.
- [76] L. Brillouin, *Wave propagation and group velocity*. Academic Press, 1960.
- [77] R. Plonsey and R. E. Collin, *Principles and applications of electromagnetic fields*. McGraw-Hill, 1961.
- [78] M. N. Sadiku, *Elements of electromagnetics*. Oxford University Press, 2014.
- [79] A. Wadhwa, A. L. Dal, and N. Malhotra, “Transmission media,” <https://www.slideshare.net/abhishekwadhw786/transmission-media-9416228>.
- [80] P. H. Smith, “Transmission line calculator,” *Electronics*, vol. 12, no. 1, pp. 29–31, 1939.
- [81] F. B. Hildebrand, *Advanced calculus for applications*. Prentice-Hall, 1962.
- [82] J. Schutt-Aine, “Experiment02-coaxial transmission line measurement using slotted line,” <http://emlab.uiuc.edu/ece451/ECE451Lab02.pdf>.
- [83] D. M. Pozar, E. J. K. Knapp, and J. B. Mead, “ECE 584 microwave engineering laboratory notebook,” [http://www.ecs.umass.edu/ece/ece584/ECE584\\_lab\\_manual.pdf](http://www.ecs.umass.edu/ece/ece584/ECE584_lab_manual.pdf), 2004.
- [84] R. E. Collin, *Field theory of guided waves*. McGraw-Hill, 1960.

- [85] Q. S. Liu, S. Sun, and W. C. Chew, "A potential-based integral equation method for low-frequency electromagnetic problems," *IEEE Transactions on Antennas and Propagation*, vol. 66, no. 3, pp. 1413–1426, 2018.
- [86] Wikipedia, "Snell's law," [https://en.wikipedia.org/wiki/Snell's\\_law](https://en.wikipedia.org/wiki/Snell's_law).
- [87] G. Tyras, *Radiation and propagation of electromagnetic waves*. Academic Press, 1969.
- [88] L. Brekhovskikh, *Waves in layered media*. Academic Press, 1980.
- [89] Scholarpedia, "Goos-hanchen effect," [http://www.scholarpedia.org/article/Goos-Hanchen\\_effect](http://www.scholarpedia.org/article/Goos-Hanchen_effect).
- [90] K. Kao and G. A. Hockham, "Dielectric-fibre surface waveguides for optical frequencies," in *Proceedings of the Institution of Electrical Engineers*, vol. 113, no. 7. IET, 1966, pp. 1151–1158.
- [91] E. Glytsis, "Slab waveguide fundamentals," [http://users.ntua.gr/eglytsis/IO/Slab-Waveguides\\_p.pdf](http://users.ntua.gr/eglytsis/IO/Slab-Waveguides_p.pdf), 2018.
- [92] Wikipedia, "Optical fiber," [https://en.wikipedia.org/wiki/Optical\\_fiber](https://en.wikipedia.org/wiki/Optical_fiber).
- [93] Atlantic Cable, "1869 indo-european cable," <https://atlantic-cable.com/Cables/1869IndoEur/index.htm>.
- [94] Wikipedia, "Submarine communications cable," [https://en.wikipedia.org/wiki/Submarine\\_communications\\_cable](https://en.wikipedia.org/wiki/Submarine_communications_cable).
- [95] D. Brewster, "On the laws which regulate the polarisation of light by reflexion from transparent bodies," *Philosophical Transactions of the Royal Society of London*, vol. 105, pp. 125–159, 1815.
- [96] Wikipedia, "Brewster's angle," [https://en.wikipedia.org/wiki/Brewster's\\_angle](https://en.wikipedia.org/wiki/Brewster's_angle).
- [97] H. Raether, "Surface plasmons on smooth surfaces," in *Surface plasmons on smooth and rough surfaces and on gratings*. Springer, 1988, pp. 4–39.
- [98] E. Kretschmann and H. Raether, "Radiative decay of non radiative surface plasmons excited by light," *Zeitschrift für Naturforschung A*, vol. 23, no. 12, pp. 2135–2136, 1968.
- [99] Wikipedia, "Surface plasmon," [https://en.wikipedia.org/wiki/Surface\\_plasmon](https://en.wikipedia.org/wiki/Surface_plasmon).
- [100] Wikimedia, "Gaussian wave packet," [https://commons.wikimedia.org/wiki/File:Gaussian\\_wave\\_packet.svg](https://commons.wikimedia.org/wiki/File:Gaussian_wave_packet.svg).
- [101] Wikipedia, "Charles K. Kao," [https://en.wikipedia.org/wiki/Charles\\_K.\\_Kao](https://en.wikipedia.org/wiki/Charles_K._Kao).
- [102] H. B. Callen and T. A. Welton, "Irreversibility and generalized noise," *Physical Review*, vol. 83, no. 1, p. 34, 1951.
- [103] R. Kubo, "The fluctuation-dissipation theorem," *Reports on progress in physics*, vol. 29, no. 1, p. 255, 1966.

- [104] C. Lee, S. Lee, and S. Chuang, "Plot of modal field distribution in rectangular and circular waveguides," *IEEE transactions on microwave theory and techniques*, vol. 33, no. 3, pp. 271–274, 1985.
- [105] W. C. Chew, *Waves and Fields in Inhomogeneous Media*. IEEE Press, 1996.
- [106] M. Abramowitz and I. A. Stegun, *Handbook of mathematical functions: with formulas, graphs, and mathematical tables*. Courier Corporation, 1965, vol. 55.
- [107] —, "Handbook of mathematical functions: with formulas, graphs, and mathematical tables," <http://people.math.sfu.ca/~cbm/aands/index.htm>.
- [108] W. C. Chew, W. Sha, and Q. I. Dai, "Green's dyadic, spectral function, local density of states, and fluctuation dissipation theorem," *arXiv preprint arXiv:1505.01586*, 2015.
- [109] Wikipedia, "Very Large Array," [https://en.wikipedia.org/wiki/Very\\_Large\\_Array](https://en.wikipedia.org/wiki/Very_Large_Array).
- [110] C. A. Balanis and E. Holzman, "Circular waveguides," *Encyclopedia of RF and Microwave Engineering*, 2005.
- [111] M. Al-Hakkak and Y. Lo, "Circular waveguides with anisotropic walls," *Electronics Letters*, vol. 6, no. 24, pp. 786–789, 1970.
- [112] Wikipedia, "Horn Antenna," [https://en.wikipedia.org/wiki/Horn\\_antenna](https://en.wikipedia.org/wiki/Horn_antenna).
- [113] P. Silvester and P. Benedek, "Microstrip discontinuity capacitances for right-angle bends, t junctions, and crossings," *IEEE Transactions on Microwave Theory and Techniques*, vol. 21, no. 5, pp. 341–346, 1973.
- [114] R. Garg and I. Bahl, "Microstrip discontinuities," *International Journal of Electronics Theoretical and Experimental*, vol. 45, no. 1, pp. 81–87, 1978.
- [115] P. Smith and E. Turner, "A bistable fabry-perot resonator," *Applied Physics Letters*, vol. 30, no. 6, pp. 280–281, 1977.
- [116] A. Yariv, *Optical electronics*. Saunders College Publ., 1991.
- [117] Wikipedia, "Klystron," <https://en.wikipedia.org/wiki/Klystron>.
- [118] —, "Magnetron," [https://en.wikipedia.org/wiki/Cavity\\_magnetron](https://en.wikipedia.org/wiki/Cavity_magnetron).
- [119] —, "Absorption Wavemeter," [https://en.wikipedia.org/wiki/Absorption\\_wavemeter](https://en.wikipedia.org/wiki/Absorption_wavemeter).
- [120] W. C. Chew, M. S. Tong, and B. Hu, "Integral equation methods for electromagnetic and elastic waves," *Synthesis Lectures on Computational Electromagnetics*, vol. 3, no. 1, pp. 1–241, 2008.
- [121] A. D. Yaghjian, "Reflections on Maxwell's treatise," *Progress In Electromagnetics Research*, vol. 149, pp. 217–249, 2014.
- [122] L. Nagel and D. Pederson, "Simulation program with integrated circuit emphasis," in *Midwest Symposium on Circuit Theory*, 1973.

- [123] S. A. Schelkunoff and H. T. Friis, *Antennas: theory and practice*. Wiley New York, 1952, vol. 639.
- [124] H. G. Schantz, “A brief history of uwb antennas,” *IEEE Aerospace and Electronic Systems Magazine*, vol. 19, no. 4, pp. 22–26, 2004.
- [125] E. Kudeki, “Fields and Waves,” <http://remote2.ece.illinois.edu/~erhan/FieldsWaves/ECE350lectures.html>.
- [126] Wikipedia, “Antenna Aperture,” [https://en.wikipedia.org/wiki/Antenna\\_aperture](https://en.wikipedia.org/wiki/Antenna_aperture).
- [127] C. A. Balanis, *Antenna theory: analysis and design*. John Wiley & Sons, 2016.
- [128] R. W. P. King, G. S. Smith, M. Owens, and T. Wu, “Antennas in matter: Fundamentals, theory, and applications,” *NASA STI/Recon Technical Report A*, vol. 81, 1981.
- [129] H. Yagi and S. Uda, “Projector of the sharpest beam of electric waves,” *Proceedings of the Imperial Academy*, vol. 2, no. 2, pp. 49–52, 1926.
- [130] Wikipedia, “Yagi-Uda Antenna,” [https://en.wikipedia.org/wiki/Yagi-Uda\\_antenna](https://en.wikipedia.org/wiki/Yagi-Uda_antenna).
- [131] Antenna-theory.com, “Slot Antenna,” <http://www.antenna-theory.com/antennas/aperture/slot.php>.
- [132] A. D. Olver and P. J. Clarricoats, *Microwave horns and feeds*. IET, 1994, vol. 39.
- [133] B. Thomas, “Design of corrugated conical horns,” *IEEE Transactions on Antennas and Propagation*, vol. 26, no. 2, pp. 367–372, 1978.
- [134] P. J. B. Clarricoats and A. D. Olver, *Corrugated horns for microwave antennas*. IET, 1984, no. 18.
- [135] P. Gibson, “The vivaldi aerial,” in *1979 9th European Microwave Conference*. IEEE, 1979, pp. 101–105.
- [136] Wikipedia, “Vivaldi Antenna,” [https://en.wikipedia.org/wiki/Vivaldi\\_antenna](https://en.wikipedia.org/wiki/Vivaldi_antenna).
- [137] —, “Cassegrain Antenna,” [https://en.wikipedia.org/wiki/Cassegrain\\_antenna](https://en.wikipedia.org/wiki/Cassegrain_antenna).
- [138] —, “Cassegrain Reflector,” [https://en.wikipedia.org/wiki/Cassegrain\\_reflector](https://en.wikipedia.org/wiki/Cassegrain_reflector).
- [139] W. A. Imbriale, S. S. Gao, and L. Boccia, *Space antenna handbook*. John Wiley & Sons, 2012.
- [140] J. A. Encinar, “Design of two-layer printed reflectarrays using patches of variable size,” *IEEE Transactions on Antennas and Propagation*, vol. 49, no. 10, pp. 1403–1410, 2001.
- [141] D.-C. Chang and M.-C. Huang, “Microstrip reflectarray antenna with offset feed,” *Electronics Letters*, vol. 28, no. 16, pp. 1489–1491, 1992.

- [142] G. Minatti, M. Faenzi, E. Martini, F. Caminita, P. De Vita, D. González-Ovejero, M. Sabbadini, and S. Maci, “Modulated metasurface antennas for space: Synthesis, analysis and realizations,” *IEEE Transactions on Antennas and Propagation*, vol. 63, no. 4, pp. 1288–1300, 2014.
- [143] X. Gao, X. Han, W.-P. Cao, H. O. Li, H. F. Ma, and T. J. Cui, “Ultrawideband and high-efficiency linear polarization converter based on double v-shaped metasurface,” *IEEE Transactions on Antennas and Propagation*, vol. 63, no. 8, pp. 3522–3530, 2015.
- [144] D. De Schweinitz and T. L. Frey Jr, “Artificial dielectric lens antenna,” Nov. 13 2001, US Patent 6,317,092.
- [145] K.-L. Wong, “Planar antennas for wireless communications,” *Microwave Journal*, vol. 46, no. 10, pp. 144–145, 2003.
- [146] H. Nakano, M. Yamazaki, and J. Yamauchi, “Electromagnetically coupled curl antenna,” *Electronics Letters*, vol. 33, no. 12, pp. 1003–1004, 1997.
- [147] K. Lee, K. Luk, K.-F. Tong, S. Shum, T. Huynh, and R. Lee, “Experimental and simulation studies of the coaxially fed U-slot rectangular patch antenna,” *IEE Proceedings-Microwaves, Antennas and Propagation*, vol. 144, no. 5, pp. 354–358, 1997.
- [148] K. Luk, C. Mak, Y. Chow, and K. Lee, “Broadband microstrip patch antenna,” *Electronics letters*, vol. 34, no. 15, pp. 1442–1443, 1998.
- [149] M. Bolic, D. Simplot-Ryl, and I. Stojmenovic, *RFID systems: research trends and challenges*. John Wiley & Sons, 2010.
- [150] D. M. Dobkin, S. M. Weigand, and N. Iyer, “Segmented magnetic antennas for near-field UHF RFID,” *Microwave Journal*, vol. 50, no. 6, p. 96, 2007.
- [151] Z. N. Chen, X. Qing, and H. L. Chung, “A universal UHF RFID reader antenna,” *IEEE transactions on microwave theory and techniques*, vol. 57, no. 5, pp. 1275–1282, 2009.
- [152] C.-T. Chen, *Linear system theory and design*. Oxford University Press, Inc., 1998.
- [153] S. H. Schot, “Eighty years of Sommerfeld’s radiation condition,” *Historia mathematica*, vol. 19, no. 4, pp. 385–401, 1992.
- [154] A. Ishimaru, *Electromagnetic wave propagation, radiation, and scattering from fundamentals to applications*. Wiley Online Library, 2017, also 1991.
- [155] A. E. H. Love, “I. the integration of the equations of propagation of electric waves,” *Philosophical Transactions of the Royal Society of London. Series A, Containing Papers of a Mathematical or Physical Character*, vol. 197, no. 287-299, pp. 1–45, 1901.
- [156] Wikipedia, “Christiaan Huygens,” [https://en.wikipedia.org/wiki/Christiaan\\_Huygens](https://en.wikipedia.org/wiki/Christiaan_Huygens).
- [157] —, “George Green (mathematician),” [https://en.wikipedia.org/wiki/George\\_Green\\_\(mathematician\)](https://en.wikipedia.org/wiki/George_Green_(mathematician)).

- [158] C.-T. Tai, *Dyadic Greens Functions in Electromagnetic Theory*. PA: International Textbook, Scranton, 1971.
- [159] —, *Dyadic Green functions in electromagnetic theory*. Institute of Electrical & Electronics Engineers (IEEE), 1994.
- [160] W. Franz, “Zur formulierung des huygensschen prinzipts,” *Zeitschrift für Naturforschung A*, vol. 3, no. 8-11, pp. 500–506, 1948.
- [161] J. A. Stratton, *Electromagnetic Theory*. McGraw-Hill Book Company, Inc., 1941.
- [162] J. D. Jackson, *Classical Electrodynamics*. John Wiley & Sons, 1962.
- [163] W. Meissner and R. Ochsenfeld, “Ein neuer effekt bei eintritt der supraleitfähigkeit,” *Naturwissenschaften*, vol. 21, no. 44, pp. 787–788, 1933.
- [164] Wikipedia, “Superconductivity,” <https://en.wikipedia.org/wiki/Superconductivity>.
- [165] D. Sievenpiper, L. Zhang, R. F. Broas, N. G. Alexopolous, and E. Yablonovitch, “High-impedance electromagnetic surfaces with a forbidden frequency band,” *IEEE Transactions on Microwave Theory and techniques*, vol. 47, no. 11, pp. 2059–2074, 1999.
- [166] Wikipedia, “Snell’s law,” [https://en.wikipedia.org/wiki/Snell's\\_law](https://en.wikipedia.org/wiki/Snell's_law).
- [167] H. Lamb, “On sommerfeld’s diffraction problem; and on reflection by a parabolic mirror,” *Proceedings of the London Mathematical Society*, vol. 2, no. 1, pp. 190–203, 1907.
- [168] W. J. Smith, *Modern optical engineering*. McGraw-Hill New York, 1966, vol. 3.
- [169] D. C. O’Shea, T. J. Suleski, A. D. Kathman, and D. W. Prather, *Diffraction optics: design, fabrication, and test*. Spie Press Bellingham, WA, 2004, vol. 62.
- [170] J. B. Keller and H. B. Keller, “Determination of reflected and transmitted fields by geometrical optics,” *JOSA*, vol. 40, no. 1, pp. 48–52, 1950.
- [171] G. A. Deschamps, “Ray techniques in electromagnetics,” *Proceedings of the IEEE*, vol. 60, no. 9, pp. 1022–1035, 1972.
- [172] R. G. Kouyoumjian and P. H. Pathak, “A uniform geometrical theory of diffraction for an edge in a perfectly conducting surface,” *Proceedings of the IEEE*, vol. 62, no. 11, pp. 1448–1461, 1974.
- [173] R. Kouyoumjian, “The geometrical theory of diffraction and its application,” in *Numerical and Asymptotic Techniques in Electromagnetics*. Springer, 1975, pp. 165–215.
- [174] S.-W. Lee and G. Deschamps, “A uniform asymptotic theory of electromagnetic diffraction by a curved wedge,” *IEEE Transactions on Antennas and Propagation*, vol. 24, no. 1, pp. 25–34, 1976.
- [175] Wikipedia, “Fermat’s principle,” [https://en.wikipedia.org/wiki/Fermat's\\_principle](https://en.wikipedia.org/wiki/Fermat's_principle).



- [176] N. Yu, P. Genevet, M. A. Kats, F. Aieta, J.-P. Tetienne, F. Capasso, and Z. Gaburro, “Light propagation with phase discontinuities: generalized laws of reflection and refraction,” *Science*, vol. 334, no. 6054, pp. 333–337, 2011.
- [177] A. Sommerfeld, *Partial differential equations in physics*. Academic Press, 1949, vol. 1.
- [178] R. Haberman, *Elementary applied partial differential equations*. Prentice Hall Englewood Cliffs, NJ, 1983, vol. 987.
- [179] G. A. Deschamps, “Gaussian beam as a bundle of complex rays,” *Electronics letters*, vol. 7, no. 23, pp. 684–685, 1971.
- [180] J. Enderlein and F. Pampaloni, “Unified operator approach for deriving hermite–gaussian and laguerre–gaussian laser modes,” *JOSA A*, vol. 21, no. 8, pp. 1553–1558, 2004.
- [181] D. L. Andrews, *Structured light and its applications: An introduction to phase-structured beams and nanoscale optical forces*. Academic Press, 2011.
- [182] J. W. Strutt, “Xv. on the light from the sky, its polarization and colour,” *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, vol. 41, no. 271, pp. 107–120, 1871.
- [183] L. Rayleigh, “X. on the electromagnetic theory of light,” *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, vol. 12, no. 73, pp. 81–101, 1881.
- [184] R. C. Wittmann, “Spherical wave operators and the translation formulas,” *IEEE Transactions on Antennas and Propagation*, vol. 36, no. 8, pp. 1078–1087, 1988.
- [185] S. Sun, Y. G. Liu, W. C. Chew, and Z. Ma, “Calderón multiplicative preconditioned efie with perturbation method,” *IEEE Transactions on Antennas and Propagation*, vol. 61, no. 1, pp. 247–255, 2012.
- [186] G. Mie, “Beiträge zur optik trüber medien, speziell kolloidaler metallösungen,” *Annalen der physik*, vol. 330, no. 3, pp. 377–445, 1908.
- [187] Wikipedia, “Mie scattering,” [https://en.wikipedia.org/wiki/Mie\\_scattering](https://en.wikipedia.org/wiki/Mie_scattering).
- [188] R. E. Collin, *Foundations for microwave engineering*. John Wiley & Sons, 2007, also 1966.
- [189] L. B. Felsen and N. Marcuvitz, *Radiation and scattering of waves*. John Wiley & Sons, 1994, also 1973, vol. 31.
- [190] P. P. Ewald, “Die berechnung optischer und elektrostatischer gitterpotentiale,” *Annalen der physik*, vol. 369, no. 3, pp. 253–287, 1921.
- [191] E. Whittaker and G. Watson, *A Course of Modern Analysis*. Cambridge Mathematical Library, 1927.

- [192] A. Sommerfeld, *Über die Ausbreitung der Wellen in der drahtlosen Telegraphie*. Verlag der Königlich Bayerischen Akademie der Wissenschaften, 1909.
- [193] J. Kong, "Electromagnetic fields due to dipole antennas over stratified anisotropic media," *Geophysics*, vol. 37, no. 6, pp. 985–996, 1972.
- [194] K. Yee, "Numerical solution of initial boundary value problems involving maxwell's equations in isotropic media," *IEEE Transactions on Antennas and Propagation*, vol. 14, no. 3, pp. 302–307, 1966.
- [195] A. Taflove, "Review of the formulation and applications of the finite-difference time-domain method for numerical modeling of electromagnetic wave interactions with arbitrary structures," *Wave Motion*, vol. 10, no. 6, pp. 547–582, 1988.
- [196] A. Taflove and S. C. Hagness, *Computational electrodynamics: the finite-difference time-domain method*. Artech house, 2005, also 1995.
- [197] W. Yu, R. Mittra, T. Su, Y. Liu, and X. Yang, *Parallel finite-difference time-domain method*. Artech House Norwood, 2006.
- [198] D. Potter, "Computational physics," 1973.
- [199] W. F. Ames, *Numerical methods for partial differential equations*. Academic press, 2014, also 1977.
- [200] K. W. Morton, *Revival: Numerical Solution Of Convection-Diffusion Problems (1996)*. CRC Press, 2019.
- [201] K. Aki and P. G. Richards, *Quantitative seismology*, 2002.
- [202] W. C. Chew, "Electromagnetic theory on a lattice," *Journal of Applied Physics*, vol. 75, no. 10, pp. 4843–4850, 1994.
- [203] J. v. Neumann, *Mathematische Grundlagen der Quantenmechanik, Berlin*. Springer, New York, Dover Publications, 1943.
- [204] R. Courant, K. Friedrichs, and H. Lewy, "Über die partiellen differenzgleichungen der mathematischen physik," *Mathematische annalen*, vol. 100, no. 1, pp. 32–74, 1928.
- [205] J.-P. Berenger, "A perfectly matched layer for the absorption of electromagnetic waves," *Journal of computational physics*, vol. 114, no. 2, pp. 185–200, 1994.
- [206] W. C. Chew and W. H. Weedon, "A 3d perfectly matched medium from modified maxwell's equations with stretched coordinates," *Microwave and optical technology letters*, vol. 7, no. 13, pp. 599–604, 1994.
- [207] W. C. Chew, J. Jin, and E. Michielssen, "Complex coordinate system as a generalized absorbing boundary condition," in *IEEE Antennas and Propagation Society International Symposium 1997. Digest*, vol. 3. IEEE, 1997, pp. 2060–2063.

- [208] W. C. H. McLean, *Strongly elliptic systems and boundary integral equations*. Cambridge University Press, 2000.
- [209] G. C. Hsiao and W. L. Wendland, *Boundary integral equations*. Springer, 2008.
- [210] K. F. Warnick, *Numerical analysis for electromagnetic integral equations*. Artech House, 2008.
- [211] M. M. Botha, “Solving the volume integral equations of electromagnetic scattering,” *Journal of Computational Physics*, vol. 218, no. 1, pp. 141–158, 2006.
- [212] P. K. Banerjee and R. Butterfield, *Boundary element methods in engineering science*. McGraw-Hill London, 1981, vol. 17.
- [213] O. C. Zienkiewicz, R. L. Taylor, P. Nithiarasu, and J. Zhu, *The finite element method*. McGraw-Hill London, 1977, vol. 3.
- [214] J.-F. Lee, R. Lee, and A. Cangelaris, “Time-domain finite-element methods,” *IEEE Transactions on Antennas and Propagation*, vol. 45, no. 3, pp. 430–442, 1997.
- [215] J. L. Volakis, A. Chatterjee, and L. C. Kempel, *Finite element method electromagnetics: antennas, microwave circuits, and scattering applications*. John Wiley & Sons, 1998, vol. 6.
- [216] J.-M. Jin, *The finite element method in electromagnetics*. John Wiley & Sons, 2015.
- [217] G. Strang, *Linear algebra and its applications*. Academic Press, 1976.
- [218] Cramer and Gabriel, *Introduction a l’analyse des lignes courbes algebriques par Gabriel Cramer...* chez les freres Cramer & Cl. Philibert, 1750.
- [219] J. A. Schouten, *Tensor analysis for physicists*. Courier Corporation, 1989.
- [220] W. H. Press, S. A. Teukolsky, W. T. Vetterling, and B. P. Flannery, *Numerical recipes 3rd edition: The art of scientific computing*. Cambridge University Press, 2007.
- [221] Wikipedia, “John Dalton,” [https://en.wikipedia.org/wiki/John\\_Dalton](https://en.wikipedia.org/wiki/John_Dalton).
- [222] —, “Max Planck,” [https://en.wikipedia.org/wiki/Max\\_Planck](https://en.wikipedia.org/wiki/Max_Planck).
- [223] —, “Photoelectric effect,” [https://en.wikipedia.org/wiki/Photoelectric\\_effect](https://en.wikipedia.org/wiki/Photoelectric_effect).
- [224] B. Bapat, “Newton’s rings,” [http://www.iiserpune.ac.in/~bhasbapat/phy221\\_files/NewtonsRing.pdf](http://www.iiserpune.ac.in/~bhasbapat/phy221_files/NewtonsRing.pdf).
- [225] Wikipedia, “Double-slit experiment,” [https://en.wikipedia.org/wiki/Double-slit\\_experiment](https://en.wikipedia.org/wiki/Double-slit_experiment).
- [226] Shmoop.Com, “Young’s double-slit,” <https://www.shmoop.com/optics/young-double-slit.html>.
- [227] Wikipedia, “Louis de Broglie,” [https://en.wikipedia.org/wiki/Louis\\_de\\_Broglie](https://en.wikipedia.org/wiki/Louis_de_Broglie).

- [228] —, “Newton’s laws of motion,” [https://en.wikipedia.org/wiki/Newton's\\_laws\\_of\\_motion](https://en.wikipedia.org/wiki/Newton's_laws_of_motion).
- [229] —, “Quantum electrodynamics,” [https://en.wikipedia.org/wiki/Quantum\\_electrodynamics](https://en.wikipedia.org/wiki/Quantum_electrodynamics).
- [230] C. Christopoulos, *The transmission-line modeling method: TLM*. IEEE New York, 1995, vol. 221.
- [231] A. E. Ruehli, G. Antonini, and L. Jiang, *Circuit oriented electromagnetic modeling using the PEEC techniques*. Wiley Online Library, 2017.
- [232] Wikipedia, “William Rowan Hamilton,” [https://en.wikipedia.org/wiki/William\\_Rowan\\_Hamilton](https://en.wikipedia.org/wiki/William_Rowan_Hamilton).
- [233] W. C. Chew, A. Y. Liu, C. Salazar-Lazaro, D. Na, and W. E. I. Sha, “Hamilton equation, commutator, and energy conservation,” *Quantum Report*, in press, 2019.
- [234] Wikipedia, “Gaussian beam,” [https://en.wikipedia.org/wiki/Gaussian\\_beam](https://en.wikipedia.org/wiki/Gaussian_beam).
- [235] M. Kira and S. W. Koch, *Semiconductor quantum optics*. Cambridge University Press, 2011.
- [236] Wikipedia, “Quantum harmonic oscillator,” [https://en.wikipedia.org/wiki/Quantum\\_harmonic\\_oscillator](https://en.wikipedia.org/wiki/Quantum_harmonic_oscillator).
- [237] —, “E.C. George Sudarshan,” [https://en.wikipedia.org/wiki/E.\\_C.\\_George\\_Sudarshan](https://en.wikipedia.org/wiki/E._C._George_Sudarshan).
- [238] —, “Roy J. Glauber,” [https://en.wikipedia.org/wiki/Roy\\_J.\\_Glauber](https://en.wikipedia.org/wiki/Roy_J._Glauber).
- [239] C. Gerry, P. Knight, and P. L. Knight, *Introductory quantum optics*. Cambridge University Press, 2005.