

EE 641 Midterm Exam  
November 4, Fall 2022

Name: \_\_\_\_\_

**Q1: Instructions (4pt)**

**Rules:** I understand that this is an open book exam that shall be done within the allotted time of 120 minutes. I can use my notes, previous posted exams and exam solutions, and web resources. However, I will not communicate with any other person other than the official exam proctors during the exam, and I will not seek or accept help from any other persons other than the official proctors.

**Signature:** \_\_\_\_\_

**Q2: MAP Estimation (35pt)**

Consider a zero mean  $N$ -dimensional GMRF,  $X$ , with a density function given by

$$p(x) = \frac{1}{z} \exp \left\{ -\frac{1}{2} x^t B x \right\} ,$$

where  $B_{i,j} = a_{(i-j) \bmod N}$  and the row/column indices have the range  $i, j \in \{0, \dots, N-1\}$ .  
(i.e., The rows and columns are indexed starting at 0.)

**Q2.1:**

What is the name given to the matrix  $B$ ?

**Q2.2:**

Give an explicit expression for the value of  $z$ .

**Q2.3:**

If for  $i \neq j$  we have that  $B_{i,j} \neq 0$ , then what do you know about the relationship between  $X_i$  and  $X_j$ ?

**Q2.4:**

Write down an expression for the conditional expectation

$$E[X_i | X_j, j \neq i] ,$$

in terms of the the function  $a_i$ .

**Q2.5:**

Write down an expression for the conditional distribution

$$p(x_i | x_j, j \neq i) ,$$

in terms of the the function  $a_i$ .

**Q3: Augmented Lagrangian and ADMM (35pt)**

Consider an inverse problem in which the map estimate is given by

$$\hat{x} = \arg \min_x \{f(x) + h(x)\}$$

where  $x \in \mathbb{R}^N$ , and the two functions,  $f : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{\infty\}$ , and  $h : \mathbb{R}^N \rightarrow \mathbb{R} \cup \{\infty\}$ , are proper closed convex functions.

**Q3.1:**

Write an equivalent expression for  $\hat{x}$  based on the constrained optimization of two variables,  $(x, v)$ .

**Q3.2:**

Write out an expression for the augmented Lagrangian  $L(x, v; u)$  which solves the problem for the proper value of  $u$ .

**Q3.3:**

Let

$$(\hat{x}, \hat{v}) = \arg \min_{(x, v)} L(x, v; u) .$$

As  $u$  is increased, what will happen to the values of  $\hat{x}$  and  $\hat{v}$ ?

**Q3.4:**

Give an algorithm for solving the problem that uses the following two functions

$$F(z) = \arg \min_x \left\{ f(x) + \frac{a}{2} \|x - z\|^2 \right\} ,$$

$$H(z) = \arg \min_x \left\{ h(x) + \frac{a}{2} \|x - z\|^2 \right\} .$$

**Q3.5:**

If we choose,

$$h(x) = \begin{cases} \infty & \text{if } \exists i \text{ s.t. } x_i < 0 \\ 0 & \text{otherwise} \end{cases} ,$$

then calculate an explicit expression for  $H(z)$ .

**Q4: Surrogate Functions** (30pt) Consider the function

$$f(x) = \frac{1}{2} \|y - Ax\|^2 ,$$

with the associated quadratic surrogate function of the form

$$f(x; x') = \frac{a}{2} \|x - x'\|^2 + b^t(x - x') + c .$$

**Q4.1:**

Calculate expressions for the gradient,  $\nabla f(x)$ , and the Hessian,  $\nabla \nabla f(x)$ .

**Q4.2:**

Calculate expressions for  $c$  and  $b$  in the surrogate function.

**Q4.3:**

Calculate the matrix  $B$  so that the following expression is a Taylor series expression of  $f(x)$  about the point  $x'$ .

$$f_T(x) = \frac{1}{2} (x - x')^t B (x - x') + b^t (x - x') + c$$

**Q4.4:**

Find an expression for  $a$  so that  $f(x; x')$  is a surrogate function for  $f(x)$ .

**Q4.5:**

Give an iterative algorithm for computing the  $\hat{x} = \arg \min_x f(x)$  by using the surrogate function.