

Advantage

④ Find the pmf/pdf of a new

$$\text{R.V } Y = f(X)$$

Similar to HW6 Q11, Q12

Ex: X is uniformly chosen from $[0, 2]$

$$Y = X^2$$

Find the pdf of X and Y .

Ans: pdf of X is straightforward


$$f_X(x) = \begin{cases} 1/2 & \text{if } 0 \leq x \leq 2 \\ 0 & \text{otherwise} \end{cases}$$

For pdf of Y , we use

pdf of $X \rightarrow$ cdf of $Y \rightarrow$ pdf of Y .

$$F_Y(y) = P(Y \leq y)$$

$$= P(X^2 \leq y) = P(-\sqrt{y} \leq X \leq \sqrt{y})$$




Three cases:


$$F_Y(y) = \begin{cases} 0 & \text{if } y < 0 \\ \int_{-\sqrt{y}}^{\sqrt{y}} f_X(x) dx & \text{if } 0 \leq y < 4 \\ \int_0^{\sqrt{y}} \frac{1}{2} dx & \text{if } 0 \leq \sqrt{y} \leq 2 \end{cases}$$

$$\int_{-\sqrt{y}}^{\sqrt{y}} f_X(x) dx \quad 4 \leq y, 2 \leq \sqrt{y}$$

$$= \int_0^2 \frac{1}{2} dx = 1$$



$$f_Y(y) = \frac{dF_Y(y)}{dy} = \begin{cases} 0 & \text{if } y < 0 \\ \frac{1}{4} \times \frac{1}{\sqrt{y}} & \text{if } 0 \leq y < 4 \\ 0 & \text{if } 4 \leq y \end{cases}$$



096

Using cdf to find new pdf is
very important. One more example:

HW6 Q12

Q: X is chosen uniformly from $(0,1)$

$$Y = \frac{-\ln(X)}{\lambda} \quad \text{for some } \lambda > 0$$


Find out the cdf & the pdf
of Y .

$$\begin{aligned} \text{Ans: } F_Y(y) &\triangleq P(Y \leq y) \\ &= P\left(\frac{-\ln(X)}{\lambda} \leq y\right) \end{aligned}$$

$$= P(-\ln(X) \geq -\lambda y)$$

$$= P(X \geq e^{-\lambda y})$$

$$= \int_{e^{-\lambda y}}^{\infty} f_X(x) dx.$$



Case 1: If $y < 0$ (then $e^{-\lambda y} > 1$)

$$F_Y(y) = 0$$

If $y \geq 0$ ($e^{-\lambda y} \leq 1$)

$$F_Y(y) = \int_{e^{-\lambda y}}^1 e^{-\lambda x} \times 1 dx = 1 - e^{-\lambda y}$$

pdt: $f_Y(y)$?

$$= \frac{d}{dy} F_Y(y) = \begin{cases} 0 & \text{if } y < 0 \\ \lambda e^{-\lambda y} & \text{if } y \geq 0. \end{cases}$$

Q: What type of R.V.s is Y ?

Ans: Y is an exponential R.V.

Q: Why this is an important question?

Ans: Computer knows how to generate a uniform R.V.s X between $(0, 1)$ ex:
"rand()" in MATLAB.

By taking $Y = -\frac{\ln(X)}{\lambda}$, the Y is an exponential R.V with more Y 's closer to zero, & Y can be extremely large



Advantage

⑤ For positive R.V.s (those X with $P(X < 0) = 0$)


$$E(X) = \int_0^\infty (1 - F_X(x)) dx$$

Ex: The $F_X(x)$ of an exponential

R.V. is $\boxed{1 - e^{-\lambda x}}$

$$\begin{aligned} \Rightarrow E(X) &= \int_0^\infty (1 - (1 - e^{-\lambda x})) dx \\ &= \int_0^\infty e^{-\lambda x} dx = \frac{1}{\lambda} \end{aligned}$$

* Generalized pdf: (Not as popular as cdf.)


Cont
pdf cdf generalized pdf Discrete


Recall $f_x(x) = \frac{d}{dx} F_x(x)$

The disadvantage is that those jumps are not differentiable.

⇒ Introduce the $\delta(x)$ impulse function

Given
Ex: $F_x(x)$
Continue from one previous example.




Q: What is the generalized pdf?

Ans: Still do differentiation, and add $\Delta \delta(x-s)$ for a Δ -jump at $x=s$.

160

$$f_X(x) = \frac{1}{2} \delta(x - \frac{1}{2}) + \begin{cases} 0 & \text{if } x < 0 \\ \frac{1}{2} & \text{if } 0 \leq x < \frac{1}{2} \\ \frac{1}{2} & \text{if } \frac{1}{2} \leq x < 1 \\ 0 & \text{if } 1 \leq x \end{cases}$$

Plot $f_X(x)$



Recall the experiment that corresponds to the above derivation:

Flip a coin, if head $X = \frac{1}{2}$
if tail, X is chosen randomly between $(0, 1)$

Q: Can we directly derive the general pdf?

Ans: With prob $\frac{1}{2}$ (head), the output will be "exactly" $\frac{1}{2}$.


(10)

\Rightarrow We need a " $\frac{1}{2}\delta(x - \frac{1}{2})$ " term for the discrete prob mass function.

② for the remaining case X is chosen uniformly from $(0, 1)$

We need $f_X(x) = \frac{1}{2} + \begin{cases} 1 & \text{if } 0 < x < 1 \\ 0 & \text{otherwise} \end{cases}$

$$\Rightarrow f_X(x) = \frac{1}{2} \delta(x - \frac{1}{2}) + \begin{cases} \frac{1}{2} & \text{if } 0 < x < 1 \\ 0 & \text{otherwise} \end{cases}$$




The result is the same as if we derive $f_X(x)$ from $F_X(x)$.

Ex: Flip a ^{fair} coin, if head, X is the outcome of a fair dice

if tail, X is chosen uniformly random from $(0, 3)$. Find the pdf of X .

Ans: $f_X(x) = \frac{1}{2} \left(\frac{1}{6} \delta(x-1) + \frac{1}{6} \delta(x-2) + \dots + \frac{1}{6} \delta(x-6) \right) + \frac{1}{2} \times \begin{cases} \frac{1}{3} & \text{if } 0 < x < 3 \\ 0 & \text{otherwise} \end{cases}$



HW6 Q10


$F_X(x)$ is a cdf of X , and
we know that

$$F_X(x) = \begin{cases} 0 & x < -\frac{\pi}{2} \\ C(1 + \sin(x)) & -\frac{\pi}{2} \leq x < \frac{\pi}{2} \\ 1 & \frac{\pi}{2} \leq x \end{cases}$$

Q1: C cannot be 1. Why?

Ans: If $C=1$.

$$F_X(x) =$$



It violates ④

Q2 If $C=\frac{1}{2}$. Show that X is a conti R.V.

$$\text{Ans: } F_X(x) =$$




is conti (no jump)

$\Rightarrow X$ is a conti R.V.


Q3: Let $C = \frac{1}{4}$, Find the generalized pdf of X

Ans: Plot $F_X(x) = 1$



$$\text{Ans: } f_X(x) = \begin{cases} \frac{1}{2} \delta(x - \frac{\pi}{2}) + 0 & \text{if } x < -\frac{\pi}{2} \\ \frac{d}{dx} \left(\frac{1}{4}(1 + \sin(x)) \right) & \text{if } -\frac{\pi}{2} \leq x < \frac{\pi}{2} \\ 0 & \text{if } \frac{\pi}{2} \leq x \end{cases}$$

$$f_X(x) =$$



Many of the existing concept can be combined. 104

* Combining generalized pdf with expectation

* In the previous example, we consider the

following Flip a ^{fair} coin, if head, X is the outcome of a fair dice

if tail, X is chosen uniformly randomly from $(0,3)$.

New question: Find $E(X)$ and $\text{Var}(X)$

$$\text{Ans: } E(X) = \int_{x=-\infty}^{\infty} x f_X(x) dx$$

$$= \int_{-\infty}^{\infty} \frac{x}{12} (\delta(x-1) + \dots + \delta(x-6)) dx$$

$$+ \int_{-\infty}^{\infty} \frac{x}{12} \left(\begin{cases} \frac{1}{3} & \text{if } 0 < x < 3 \\ 0 & \text{otherwise} \end{cases} \right) dx$$

$$= \frac{1}{12} (1 + 2 + 3 + 4 + 5 + 6)$$

$$+ \int_0^3 \frac{x}{6} dx \quad \therefore \int_{-\infty}^{\infty} x \delta(x-a) dx = a$$

$$= \frac{7}{4} + \frac{3}{4} = \frac{10}{4} = \frac{5}{2}$$

$\text{Var}(X)$. exercise

105

Combining conditional prob with cdf.

* Conditional cdf

Definition:

$$F_X(x | a \leq X \leq b)$$

$$\triangleq P(X \leq x | a \leq X \leq b)$$

$$= \frac{P(X \leq x \text{ and } a \leq X \leq b)}{P(a \leq X \leq b)}$$

Example: X is a fair dice

What is the conditional cdf

given $0.3 \leq X \leq 4.0$.

Ans: $F_X(x) = P(X \leq x | 0.3 \leq X \leq 4.0)$

$$= \frac{P(X \leq x \text{ and } 0.3 \leq X \leq 4.0)}{P(0.3 \leq X \leq 4.0)}$$

$$= \begin{cases} 0 & \text{if } x < 1 \\ \frac{\frac{1}{6}}{\frac{4}{6}} = \frac{1}{4} & \text{if } 1 \leq x < 2 \\ \dots & \dots \\ \frac{\frac{1}{6}}{\frac{4}{6}} = \frac{1}{4} & \text{if } 3 \leq x < 4 \\ \frac{1}{6} & \text{if } 4 \leq x \end{cases}$$

$2 \leq x < 3$

$3 \leq x < 4$

$4 \leq x$

lot

Conditional pmf:

$$P(X=k \mid a \leq X \leq b) = \frac{P_k}{\sum_{a \leq s \leq b} P_s}$$

Conditional pdf: $f_X(x \mid a \leq X \leq b)$

Solution 1: differentiate the conditional cdf

$$f_X(x \mid a \leq X \leq b) = \frac{d}{dx} F_X(x \mid a \leq X \leq b)$$

Solution 2:

$$f_X(x \mid a \leq X \leq b) = \begin{cases} 0 & \text{if } x \text{ is not between } a \leq x \leq b. \\ c \cdot f_X(x) & \text{if } x \text{ is in } a \leq x \leq b \end{cases}$$

Where $f_X(x)$ is the original pdf.

We just need to compute the normalization coefficient c & make sure

$$\int_a^b c f_X(x) dx = 1$$

$$\Leftrightarrow C = \frac{1}{\int_a^b f_X(s) ds}$$

* Conditional Expectation / Variance

107

Note Title

2/23/2011

Combination of conditional prob,
pmf/pdf, and expectation

Ex: X is a unfair dice. with W.A
 $\begin{array}{c} 1 \\ \frac{1}{7} \\ 2 \\ \frac{1}{7} \\ 3 \\ \frac{1}{7} \\ 4 \\ \frac{1}{7} \\ 5 \\ \frac{1}{7} \\ 6 \\ \frac{1}{7} \end{array}$
Conditioning on $0.3 \leq X \leq 4.0$

What is the conditional expectation
of X , What is the conditional
variance

Ans: Step 1: Find the conditional
pdf/pmf

(Sometimes, you may need
to start from the ^{conditional} cdf)

Step 2: Compute the weighted
average using the conditional
pmf/pdf.

Step 1: $P(X=k | 0.3 \leq X \leq 4.0)$

$$= \begin{cases} 0 & \text{if } k \leq 0 \text{ or } k \geq 5 \\ \frac{P_k}{P_1 + P_2 + P_3 + P_4} & \text{if } 1 \leq k \leq 4 \end{cases}$$

$$= \begin{cases} \frac{2}{5} & \text{if } k=1 \\ \frac{1}{5} & \text{if } k=2, 3, 4 \\ 0 & \text{otherwise} \end{cases}$$

Step 2: Conditional expectation

$$E(X \mid 0.3 \leq X \leq 4.0)$$

\Rightarrow conditioning on

$$\approx 1 \times \frac{2}{5} + 2 \times \frac{1}{5} + 3 \times \frac{1}{5} + 4 \times \frac{1}{5}$$

$$\approx \frac{11}{5} *$$

$$\text{Var}(X \mid 0.3 \leq X \leq 4.0)$$

$$= E(X^2 \mid 0.3 \leq X \leq 4.0) - \left(\frac{11}{5}\right)^2$$

$$= 1^2 \times \frac{2}{5} + 2^2 \times \frac{1}{5} + 3^2 \times \frac{1}{5} + 4^2 \times \frac{1}{5}$$

$$- \left(\frac{11}{5}\right)^2$$

$$= \left(\frac{31}{5}\right) - \left(\frac{11}{5}\right)^2 = \frac{34}{25}$$

Other (unifying) descriptions of a R.V.

① Characteristic function of a R.V.

Why so many ways to describe a W.A.? Mathematicians are hoping that by using a different way of describing the W.A., the counting part can be easier

- ② Moment generating function of a R.V.
- ③ Prob. generating function
- ④ Cdf,
- ⑤ generalized pdf?

* Characteristic function of a R.V.

$\Phi_X(w)$ is a function of a parameter w

$$\widehat{\Phi}_X(w) \triangleq E(e^{jwX})$$

Ex: X is a Bernoulli R.V with para
 $p = \frac{1}{\pi}$

Find the characteristic function

$$\underline{\Phi}_X(w)$$

$$\text{Ans: } \underline{\Phi}_X(w) = E(e^{jwX})$$

$$\begin{aligned}
 &= e^{jw \cdot 0} \cdot \left(1 - \frac{1}{\pi}\right) + e^{jw \cdot 1} \left(\frac{1}{\pi}\right) \\
 &= \left(1 - \frac{1}{\pi}\right) + \frac{1}{\pi} e^{jw}
 \end{aligned}$$

Ex: X is a binomial R.V w. para

$$n=20 \quad p=0.7$$

Find $\Phi_X(w)$.

Before solving $\Phi_X(w)$, We need the binomial theorem

$$(P. 61) \quad (a+b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{(n-k)} \quad \begin{array}{l} \text{ex: } a=\pi \\ \text{b}=0.3 \end{array}$$

Why \hookrightarrow holds?

$$\therefore \text{We know } I = \sum_{k=0}^{\infty} \binom{n}{k} p^k (1-p)^{n-k}$$

for all $0 < p < 1$.

$$\text{Let } p = \frac{a}{a+b}$$

$$\therefore I = \sum_{k=0}^{\infty} \binom{n}{k} \left(\frac{a}{a+b}\right)^k \left(\frac{b}{a+b}\right)^{n-k}$$

$$\therefore (a+b)^n = \sum_{k=0}^{\infty} \binom{n}{k} a^k b^{n-k}$$

$$\text{Ans: } \Phi_X(w) = E(e^{jwX})$$

$$= \sum_{k=0}^n e^{jw k} \binom{n}{k} p^k (1-p)^{n-k}$$

III

$$= \sum_{k=0}^n \binom{n}{k} \underbrace{(e^{j\omega} p)^k}_{a} \underbrace{(-p)^{n-k}}_{b}$$

$$= (a+b)^n = (e^{j\omega} p + (-p))^n$$

What if X is a anti R.V.

$$\Phi_X(\omega) = E(e^{j\omega X})$$

$$= \int_{-\infty}^{\infty} e^{j\omega x} f_x(x) dx$$

$$= \int_{-\infty}^{\infty} f_x(x) e^{j\omega x} dx$$

Is similar to the Fourier transform of $f_x(\omega)$

$$\Rightarrow f_x(x) = \frac{1}{2\pi} \int_{\omega=-\infty}^{\infty} \Phi_X(\omega) e^{-j\omega x} d\omega$$

Ex: X is exponential R.V with para

Find $\Phi_X(\omega)$?

Ans: $\Phi_X(w) = E(e^{jwX})$

$$\begin{aligned} &= \int_{-\infty}^{\infty} e^{jwX} f_X(x) dx \\ &= \int_0^{\infty} e^{jwX} \lambda e^{-\lambda x} dx \\ &= \frac{\lambda}{\lambda - jw} \end{aligned}$$

How to use $\Phi_X(w)$?

Ans: One way of using $\Phi_X(w)$ is

the moment theorem

* For any discrete/conti/mixed type R.V
 X with $\Phi_X(w)$

$E(X^n)$ the n-th moment

$$= \left(\frac{1}{j^n} \right) \left[\frac{d^n}{dw^n} \Phi_X(w) \right]_{w=0}$$

Step 1: differentiate it n times.

Step 2: then evaluate the value
 by plugging $w=0$.

Ex: X has $\Phi_X(w) = \frac{2}{2 - jw}$

Q: What type of R.Vs is X ?

Ans: X is exponential with

para $\lambda \Rightarrow$ Now we have
① pdf/pmf ② cdf

③ Characteristic function

Q: $E(X) = ?$

$$\text{Ans: } E(X) = \frac{1}{j} \left(\frac{d}{dw} \frac{2}{2 - jw} \right) \Big|_{w=0} \quad \begin{array}{l} \text{to describe} \\ \text{a R.V.} \end{array}$$

$$= \frac{1}{j} \times \frac{2}{(2 - jw)^2} \times (-j) \Big|_{w=0}$$

$$= \frac{1}{j} \times \frac{2j}{(2 - jw)^2} \Big|_{w=0} = \frac{1}{j^2} = \frac{1}{\lambda}$$

Q: $E(X^2) = ?$

$$\text{A: } E(X^2) = \frac{1}{j^2} \left(\frac{d^2}{dw^2} \Phi_X(w) \right) \Big|_{w=0}$$

$$= \frac{1}{j^2} \left(\frac{d}{dw} \frac{2j}{(2 - jw)^2} \right) \Big|_{w=0}$$

$$= \frac{1}{j^2} \left((-2) \frac{2j}{(2 - jw)^3} \times (-j) \right) \Big|_{w=0}$$