


A Decision Model for Psychophysics (*Cont.*)

Reading: Macmillan & Creelman, Chaps. 1 & 2


Decision Model for 1-I Exp.



■ A (Perceptual) Decision Space

- ◆ x : random variable (“decision axis”)
- ◆ Each stimulus presentation determines a value of x
- ◆ $p(x|S)$: conditional probability density function
- ◆ M : mean/expected value


$$M = \int_{-\infty}^{+\infty} x p(x | S) dx$$



$$F = P(R_2 | S_1) = \int_k^{\infty} p(x | S_1) dx$$

$$H = P(R_2 | S_2) = \int_k^{\infty} p(x | S_2) dx$$


Gaussian Model of Density Functions (assuming equal variance)



$$p(x|S_1) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-M_1)^2}{2\sigma^2}}$$

$$p(x|S_2) = \frac{1}{\sqrt{2\pi}\sigma} e^{-\frac{(x-M_2)^2}{2\sigma^2}}$$


Sensitivity Index d'



$$d' = \frac{M_2 - M_1}{\sigma}$$

- d' is the normalized distance between means
- $d'=1$ defines the threshold
- d' does not depend on k

Response Bias c



$$c = \frac{1}{\sigma} \left(k - \frac{M_1 + M_2}{2} \right)$$

- c is the normalized distance between k and *average of means*
- Point of zero bias
- c can change independently of d' (How?)

How to Compute d' and c from Experimental Data?

$$\{H, F\} \longrightarrow \{z(H), z(F)\} \longrightarrow \{d', c\}$$

	R_1	R_2
S_1	$f(R_1/S_1)$	$F=f(R_2/S_1)$
S_2	$f(R_1/S_2)$	$H=f(R_2/S_2)$

Normal Deviates

$$H = \int_{-\infty}^{z(H)} \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx$$

$$F = \int_{-\infty}^{z(F)} \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx$$

$$d' = z(H) - z(F)$$


$$c = -\frac{z(H) + z(F)}{2}$$

ROC for d'

- **ROC: Receiver Operating Characteristic (Isosensitivity Curve)**
- **Question: Given the same pair of S_1 and S_2 (d' is fixed), how would performance (H , F) vary with k ?**
- **ROC plotted as $z(H)$ vs. $z(F)$ has a particularly simple form:**

$$z(H) = z(F) + d'$$

ROC: $z(\mathbf{H}) = z(\mathbf{F}) + d'$




ROC for c

- **Isobias Curve**
- **Question: Given the same criterion (c is fixed), how would performance (H, F) vary with S_1 and/or S_2 ?**
- **ROC plotted as $z(H)$ vs. $z(F)$ has a particularly simple form:**

$$z(H) = -z(F) - 2c$$


$$\text{ROC: } z(\text{H}) = -z(\text{F}) - 2c$$



How to Design a 1-I Exp.?


- How to choose S_1 and S_2 ?
- How to choose probabilities of presenting S_1 or S_2 on each trial?
- The issue of Stimulus-Response compatibility (S-R compatibility)
- How many trials?
- How many subjects?
- How to detect “bad” subjects?

More on Sensitivity Index d'



- Possible values: $0 - 4.65$ ($H=.99$, $F=.01$)
- Avoid $d'=0.0$ and $d'=\infty$
- My preference: $0.5 - 2.5$
- My preference: keep $c < 10\% \times d'$

More on Response Bias c



- If c is large, investigate why (unless it is part of the experimental design).

Your Results

- **Do you understand how the experimental data are organized, and how the different results are calculated?**
- **Do you think the results are as expected?**